

PUSH BACK ON CHRONIC HEART FAILURE

SPEND LESS  
TIME HERE...

Corlanor®  
(ivabradine) 5 mg  
7.5 mg tablets

Corlanor® was proven to reduce the chance of hospitalization from chronic heart failure.

...MORE  
TIME HERE

Corlanor® is a prescription medicine used:

- to treat adults who have chronic (lasting a long time) heart failure, with symptoms, to reduce their risk of hospitalization for worsening heart failure.

## IMPORTANT SAFETY INFORMATION

### Who should not take Corlanor®?

Do not take Corlanor® if you have:

- symptoms of heart failure that recently worsened
- very low blood pressure (hypotension)
- certain heart conditions: sick sinus syndrome, sinoatrial block, or 3rd-degree atrioventricular block
- a slow resting heart rate before treatment with Corlanor®. Ask your doctor what a slow resting heart rate is for you.

- certain liver problems
- been prescribed any medicines that can increase the effects of Corlanor®.

Ask your doctor if you are not sure if you have any of the medical conditions listed above.

Please see the following pages for additional Important Safety Information

For more information, visit [www.Corlanor.com](http://www.Corlanor.com)

## Corlanor® (ivabradine) Indication

Corlanor® is a prescription medicine used:

- to treat adults who have chronic (lasting a long time) heart failure, with symptoms, to reduce their risk of hospitalization for worsening heart failure.

## IMPORTANT SAFETY INFORMATION

### What is the most important information I should know about Corlanor®?

Corlanor® may cause serious side effects, including:

- **Harm to an unborn baby.** Females who are able to get pregnant must use effective birth control during treatment with Corlanor®. Tell your doctor right away if you become pregnant during treatment with Corlanor®.
- **Increased risk of irregular or rapid heartbeat (atrial fibrillation or heart rhythm problems).** Tell your doctor if you feel any of the following symptoms of an irregular or rapid heartbeat: heart is pounding or racing (palpitations), chest pressure, worsened shortness of breath, near fainting or fainting.
- **Slower than normal heart rate (bradycardia).** Tell your doctor if you have a slowing of heart rate, **or** symptoms of a slow heart rate such as dizziness, fatigue, lack of energy.

### Who should not take Corlanor®?

**Do not** take Corlanor® if you have:

- symptoms of heart failure that recently worsened
- very low blood pressure (hypotension)
- certain heart conditions: sick sinus syndrome, sinoatrial block, or 3rd-degree atrioventricular block
- a slow resting heart rate before treatment with Corlanor®. Ask your doctor what a slow resting heart rate is for you.
- certain liver problems
- been prescribed any medicines that can increase the effects of Corlanor®.

Ask your doctor if you are not sure if you have any of the medical conditions listed above.

### What should I tell my doctor before taking Corlanor®?

**Before you take Corlanor®, tell your doctor about all of your medical conditions, including if you:**

- have any other heart problems, including heart rhythm problems, a slow heart rate, or a heart conduction problem.
- are breastfeeding or planning to breastfeed. It is not known if Corlanor® passes into breast milk. You and your doctor should decide if you will take Corlanor® or breastfeed; do not do both.
- are pregnant or planning to become pregnant.  
**See “What is the most important information I should know about Corlanor®? – Harm to an unborn baby” section.**

Tell your doctor about all the medicines you take, including prescription and over the counter medicines, vitamins, and herbal supplements. Corlanor® may affect the way other medicines work, and other medicines may affect how Corlanor® works. This could cause serious side effects.

### What are the possible side effects of Corlanor®?

**Corlanor® may cause serious side effects. See “What is the most important information I should know about Corlanor®?”**

The most common side effects of Corlanor® are:

- increased blood pressure
- temporary brightness in part of your field of vision. This is usually caused by sudden changes in light (luminous phenomena). This brightness usually happens within the first 2 months of treatment with Corlanor® and may go away during or after treatment with Corlanor®. Be careful when driving or operating machinery where sudden changes in light can happen, especially when driving at night.

These are not all the side effects of Corlanor®. Ask your doctor or pharmacist for more information. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

**[Please click here for Full Prescribing Information and Medication Guide.](#)**

# Corlanor® is a medicine that can help treat your chronic heart failure.

**When added to beta-blockers, Corlanor® has been proven to reduce the risk of hospitalization from chronic heart failure.**

Corlanor® is a medicine that you take in addition to your current treatments for chronic heart failure.

Corlanor® is intended for certain people with chronic heart failure who have the following issues:

- Lower-left part of the heart can't pump enough blood
- Have heart failure symptoms that are stable
- A normal heart rate of 70 beats in a minute or higher at rest
- Taking beta-blockers at the highest dose they can tolerate or have a reason they cannot take beta-blockers



## How **Corlanor®** **WORKS**

Corlanor® can help lower heart rate so your heart doesn't have to work as hard.



## How **Corlanor®** **CAN HELP**

When added to your other chronic heart failure medications, Corlanor® can reduce your risk of going to the hospital due to worsening heart failure.



## How **Corlanor®** **IS TAKEN**

Corlanor® is a tablet you take two times a day with food, such as in the morning with breakfast and with your evening meal. Always follow your doctor's specific instructions, and take all of your medications as directed.

**Find out if you can save on prescription costs with the Corlanor® Copay Card.**

*See details on the following page ▶*

For more information, visit [www.Corlanor.com](http://www.Corlanor.com)

**Corlanor®**  
(ivabradine) <sup>5mg</sup>/<sub>7.5mg</sub> tablets

# Corlanor® Support

Support Services for Providers & Patients

Call **1-844-6CORLANOR** (1-844-626-7526)

for more information on the following programs and services.

## Corlanor® Copay Card\*

**Pay no more than \$20  
for each prescription\*\***

\*\*For patients with commercial insurance, the Corlanor® Copay Card may pay up to \$160 per month toward your prescription cost after you pay no more than \$20 per month for Corlanor®.



Visit [www.Corlanor.com/corlanor-copay-card/](http://www.Corlanor.com/corlanor-copay-card/) to find out if you qualify for the Corlanor® Copay Card.

\*The Corlanor® Copay Card is not available for patients receiving prescription reimbursement under any federal, state, or government-funded healthcare programs such as Medicare, Medicare Advantage, Medicare Part D, or Medicaid.

## Additional Financial Support



If you are uninsured and having difficulty paying for Corlanor®, the Amgen Safety Net Foundation may be able to help. To see if you meet the eligibility criteria please visit [amgensafetynetfoundation.com](http://amgensafetynetfoundation.com) to complete an eligibility application.



## Questions?

Call 1-844-6CORLANOR  
(1-844-626-7526)  
Monday - Friday 9AM - 9PM ET

Visit [www.Corlanor.com](http://www.Corlanor.com) to learn more about:

- Chronic heart failure causes and symptoms
- Treatments and lifestyle changes
- How Corlanor® works to help you push back
- Support for your treatment
- Advice for friends and family
- Corlanor® Copay Card and other financial help

Please click here for [Full Prescribing Information and Medication Guide](#).

**AMGEN®**

Cardiovascular